

Antimicrobial Susceptibility Testing

MICRONAUT-S Anaerobes MIC

True MIC values by broth microdilution

Innovation with Integrity

What can it be used for?

The increasing microbial resistance observed for anaerobic bacteria requires a standardized routine antimicrobial susceptibility test (AST) of anaerobes in the microbiological lab. Based on the broth microdilution (BMD) procedure, the AST plate MICRONAUT-S Anaerobes MIC provides the tool for phenotypic detection of clinically relevant resistance mechanisms in anaerobic pathogens such as *Clostridium* spp., *Bacteroides* spp. and *Peptostreptococcus* spp.

Antibiotics

The plate configuration includes antibiotics such as metronidazole, carbapenems, beta-lactamase inhibitor combinations, tigecycline or moxifloxacin, showing good antimicrobial efficacy against anaerobic bacteria.

- Amoxicillin/Clavulanate
- Ampicillin
- Clindamycin
- Doxycycline
- Ertapenem
- Imipenem
- Meropenem

- Metronidazole
- Moxifloxacin
- Penicillin G
- Piperacillin/ Tazobactam
- Tigecycline
- Vancomycin

Features and benefits

- Standardized MIC determination of antibiotics with MICRONAUT-Wilkins-Chalgren Broth in routine laboratory diagnostics
- Due to the spectrum of antibiotics, the MIC plate is appropriate for testing anaerobic bacteria originating from hospital and community acquired infections
- Reading and evaluation of the AST results can be performed visually or photometrically
- The MICRONAUT software provides reading, evaluation and interpretation according to EUCAST or CLSI criteria

Shelf life and storage

- MICRONAUT-S Anaerobes MIC, Shelf life: 24 months from date of production, Storage: at room temperature (15-25°C)
- MICRONAUT-Wilkins-Chalgren Broth, Shelf life: 12 months from date of production, Storage: at 2-8°C

Procedure

- Prepare a 0.5 McFarland standard bacteria suspension in NaCl
- Transfer into MICRONAUT-Wilkins-Chalgren Broth
- Inoculate the MICRONAUT-S Anaerobes MIC plate
- Incubate for 24-48 hours at 35-37°C under anaerobic atmosphere
- Read the result visually or measure photometrically

Antibiotics & Concentrations (µg/mL)

Amoxicillin/Clavulanate	64/32	32/16	16/8	8/4	4/2	2/1	1/0.5	0.5/0.25
Ampicillin	8	4	2	1	0.5	0.25	0.125	0.0625
Clindamycin	8	4	2	1	0.5	0.25	0.125	0.0625
Doxycycline	16	8	4	2	1	0.5	0.25	0.125
Ertapenem	16	8	4	2	1	0.5	0.25	0.125
Imipenem	64	32	16	8	4	2	1	0.5
Meropenem	64	32	16	8	4	2	1	0.5
Metronidazole	32	16	8	4	2	1	0.5	0.25
Moxifloxacin	8	4	2	1	0.5	0.25	0.125	0.0625
Penicillin G	8	4	2	1	0.5	0.25	0.125	0.0625
Piperacillin/Tazobactam	128/4	64/4	32/4	16/4	8/4	4/4	2/4	1/4
Tigecycline	8	4	2	1	-	-	-	-
Vancomycin	8	4	2	-	-	-	-	-

Order information

MICRONAUT-S Anaerobes MIC

1 test per plate, 40 plates per box / Part No. E1-085-040

MICRONAUT-Wilkins-Chalgren Broth

20 tubes per box / Part No. E2-330-020

Please contact your local representative for availability in your country. Not for sale in the USA.





Online information bruker.com/microbiology



Bruker Daltonics GmbH and Co. KG

Bremen · Germany Phone +49 (0) 421-2205-0





No. 1812818 @ 02-2023 Bruker Daltonics