



MALDI Biotyper®

# MBT HT Mycobacteria Module

---

Confident mycobacteria identification

Innovation with Integrity

CE-IVD | RUO

# Confident mycobacteria identification

The genus *Mycobacterium* includes, as major groups, the important pathogens of the *Mycobacterium tuberculosis* complex (MTC) and the nontuberculous mycobacteria (NTM). Members of the latter group are cited increasingly as the cause for opportunistic infections among immunocompromised patients. This trend and the rise of antibiotic resistance in this genus demand improved differentiation among *Mycobacterium* species.



## Covering the vast majority of Mycobacteria

The MBT HT Mycobacteria Module is the comprehensive solution for laboratories in need of highly reliable and fast mycobacteria identification via MALDI-TOF mass spectrometry. It is composed of a software module and a specific reference spectrum library covering most of the currently known mycobacteria species.

The accompanying software, which is part of the MBT HT Mycobacteria Module, triggers adapted data acquisition and analysis of mycobacteria samples, securing highly sensitive identifications.

## Optimized and safe extraction

In general, *Mycobacterium* species are more demanding for MALDI-TOF MS analysis than most other bacteria. This is due to the rigid cell wall and a low metabolic activity, which is associated with a lower number of ribosomes.

As ribosomal proteins are the main analyte molecules for micro-organism identification by MALDI-TOF, an optimized sample preparation protocol is key to yielding good quality spectra.


*Mycobacterium* samples need to be considered as biosafety level 3 organisms until the identification result is available, hence reliable inactivation is an indispensable step of the sample preparation protocol.

The above needs were addressed while developing the MBT Mycobacteria Kit, which offers a dedicated and safe sample preparation method for *Mycobacterium* spp. The kit facilitates mycobacteria sample preparation for samples cultivated in liquid as well as on solid media.

# MBT Mycobacteria Kit

## One workflow for solid media and liquid cultures

The new MBT Mycobacteria Kit offers a dedicated sample preparation method for *Mycobacterium* spp. cultivated in liquid as well as on solid media. Unlike other procedures, the inactivation method of this kit does not require boiling but comprises a convenient and user-friendly incubation of the bacteria with an Inactivation Reagent, at room temperature.



MBT Mycobacteria Kit



# MBT HT Mycobacteria Module


## Dedicated software

As *Mycobacterium* species are generally more demanding for MALDI-TOF MS analysis than most other bacteria, the comprehensive library is accompanied by a carefully designed software module.

Besides optimized sample preparation, also adapted data acquisition and subsequent data analysis are crucial to yielding good identification results. Thorough validation of the matching scores resulted in reliably accepting lower thresholds for *Mycobacterium* spp., ensuring highly specific and sensitive identification.

Selection of the appropriate sample type as “Mycobacteria” while setting up the analysis run automatically triggers the use of the dedicated software module for data acquisition, followed by matching the unknown spectrum against the reference spectra incorporated in the MBT HT Mycobacteria Module, taking into account the specific log(score) thresholds for reliable identification.

Sample	
<input type="checkbox"/>	BTS
<input type="checkbox"/>	Filamentous Fungi
<input type="checkbox"/>	Mycobacteria
<input type="checkbox"/>	Sepsityper



The screenshot shows the 'Sample type' selection interface. A dropdown menu is open, showing options: 'Sample', 'BTS', 'Filamentous Fungi', 'Mycobacteria', 'Sepsityper', and 'Sample'. The 'Mycobacteria' option is highlighted. The background shows a grid of sample wells and a table with columns: 'Position', 'Sample Identifier', 'Name', 'Sample type', 'Preparation protocol', 'Isolate Identifier', and 'Description'.

Easy selection of the sample type while setting up the run for optimized data acquisition and analysis

## One reference library for solid media and liquid cultures

Spectra of most mycobacteria strains grown on solid Löwenstein-Jensen or BACTEC™ MGIT™ (BD Diagnostics) liquid media show no significant variation. For those with varying spectra, reference spectra from both liquid and solid media are stored in the mycobacteria reference library.

## International mycobacteria consortium

The aim of this consortium is to collect securely identified mycobacteria strains to create a library as the basis for the highest performance in mycobacteria identification.

# Library content

## 182 species entries

<i>M. abscessus</i>	<i>M. diernhoferi</i>	<i>M. kyorinense</i>	<i>M. pseudoshottsii</i>
<i>M. africanum</i>	<i>M. doricum</i>	<i>M. lacus</i>	<i>M. psychrotolerans</i>
<i>M. agri</i>	<i>M. duvalii</i>	<i>M. lehmannii</i>	<i>M. pulveris</i>
<i>M. aichiense</i>	<i>M. eburneum</i>	<i>M. lentiflavum</i>	<i>M. pyrenivorans</i>
<i>M. algericum</i>	<i>M. elephantis</i>	<i>M. litorale</i>	<i>M. rhodesiae</i>
<i>M. alsense</i>	<i>M. engbaekii</i>	<i>M. llatzerense</i>	<i>M. riyadhense</i>
<i>M. alvei</i>	<i>M. europaeum</i>	<i>M. longobardum</i>	<i>M. rufum</i>
<i>M. angelicum</i>	<i>M. fallax</i>	<i>M. lutetiense</i>	<i>M. rutilum</i>
<i>M. anyangense</i>	<i>M. farcinogenes</i>	<i>M. madagascariense</i>	<i>M. salmoniphilum</i>
<i>M. aquaticum</i>	<i>M. flavescens</i>	<i>M. mageritense</i>	<i>M. saopaulense</i>
<i>M. arabiense</i>	<i>M. florentinum</i>	<i>M. malmoense</i>	<i>M. sarraceniae</i>
<i>M. arcueilense</i>	<i>M. fluoranthenivorans</i>	<i>M. mantenii</i>	<i>M. saskatchewanense</i>
<i>M. aromaticivorans</i>	<i>M. fortuitum</i>	<i>M. marinum</i>	<i>M. scrofulaceum</i>
<i>M. arosiense</i>	<i>M. fragae</i>	<i>M. marseillense</i>	<i>M. sediminis</i>
<i>M. arupense</i>	<i>M. franklinii</i>	<i>M. microti</i>	<i>M. senegalense</i>
<i>M. asiaticum</i>	<i>M. frederiksbergense</i>	<i>M. minnesotense</i>	<i>M. senuense</i>
<i>M. aubagnense</i>	<i>M. gadium</i>	<i>M. monacense</i>	<i>M. seoulense</i>
<i>M. aurum</i>	<i>M. gastris</i>	<i>M. montefiorensis</i>	<i>M. septicum</i>
<i>M. austroafricanum</i>	<i>M. genavense</i>	<i>M. montmartrensis</i>	<i>M. setense</i>
<i>M. avium</i>	<i>M. gilvum</i>	<i>M. moriokaense</i>	<i>M. sherrisii</i>
<i>M. bacteremicum</i>	<i>M. goodii</i>	<i>M. mucogenicum</i>	<i>M. shigaense</i>
<i>M. basiliense</i>	<i>M. gordonae</i>	<i>M. murale</i>	<i>M. shimoidei</i>
<i>M. boenickei</i>	<i>M. grossiae</i>	<i>M. nebraskense</i>	<i>M. shinjukuense</i>
<i>M. bohemicum</i>	<i>M. haemophilum</i>	<i>M. neoaurum</i>	<i>M. simiae</i>
<i>M. botniense</i>	<i>M. hassiacum</i>	<i>M. neumannii</i>	<i>M. smegmatis</i>
<i>M. bovis</i>	<i>M. heckeshornense</i>	<i>M. neworleansense</i>	<i>M. sphagni</i>
<i>M. bourgelatii</i>	<i>M. heidelbergense</i>	<i>M. nonchromogenicum</i>	<i>M. stephanolepidis</i>
<i>M. branderi</i>	<i>M. helvum</i>	<i>M. noviomagense</i>	<i>M. stomatepiae</i>
<i>M. brisbanense</i>	<i>M. heraklionense</i>	<i>M. novocastrensis</i>	<i>M. szulgai</i>
<i>M. brumae</i>	<i>M. hiberniae</i>	<i>M. obuense</i>	<i>M. talmoniae</i>
<i>M. canariensis</i>	<i>M. hippocampi</i>	<i>M. pallens</i>	<i>M. terrae</i>
<i>M. caprae</i>	<i>M. hodleri</i>	<i>M. palustre</i>	<i>M. thermoresistibile</i>
<i>M. celatum</i>	<i>M. holsaticum</i>	<i>M. paraense</i>	<i>M. timonense</i>
<i>M. celeriflavum</i>	<i>M. houstonense</i>	<i>M. paraffinicum</i>	<i>M. tokaiense</i>
<i>M. chelonae</i>	<i>M. icosiumassiliensis</i>	<i>M. parafortuitum</i>	<i>M. triplex</i>
<i>M. chimaera</i>	<i>M. immunogenum</i>	<i>M. paragordonae</i>	<i>M. triviale</i>
<i>M. chitae</i>	<i>M. insubricum</i>	<i>M. parakoreense</i>	<i>M. tuberculosis</i>
<i>M. chlorophenolicum</i>	<i>M. interjectum</i>	<i>M. parascrofulaceum</i>	<i>M. tusciae</i>
<i>M. chubuense</i>	<i>M. intermedium</i>	<i>M. paraseoulense</i>	<i>M. vaccae</i>
<i>M. colombiense</i>	<i>M. intracellulare</i>	<i>M. paraterrae</i>	<i>M. vanbaalenii</i>
<i>M. conceptionense</i>	<i>M. iranicum</i>	<i>M. parmense</i>	<i>M. virginense</i>
<i>M. confluentis</i>	<i>M. kansasii</i>	<i>M. peregrinum</i>	<i>M. vulnerans</i>
<i>M. conspicuum</i>	<i>M. komossense</i>	<i>M. phlei</i>	<i>M. wolinskyi</i>
<i>M. cookii</i>	<i>M. koreense</i>	<i>M. phocaicum</i>	<i>M. xenopi</i>
<i>M. cosmeticum</i>	<i>M. kubicae</i>	<i>M. porcinum</i>	
<i>M. crocinum</i>	<i>M. kumamotoense</i>	<i>M. poriferae</i>	

## Order information



### **MBT HT Mycobacteria IVD Module**

Part No. 1877012

Consists of the MBT HT Mycobacteria IVD Module Library and license for the dedicated software module. Prerequisite for the module is the MBT Compass HT IVD software.



### **MBT Mycobacteria IVD Kit**

Part No. 1889166

Containing all reagents and consumables required for mycobacteria preparation of samples cultivated on solid and in liquid media.

Please contact your local representative for availability in your country. Not for sale in the USA.



### **MBT HT Mycobacteria Module**

Part-No. 1889530

Consists of the MBT Mycobacteria Library and license for the dedicated software module. Prerequisite for the module is the MBT Compass HT software.

### **MBT Mycobacteria Kit**

Part-No. 1889119

Containing all reagents and consumables required for mycobacteria preparation of samples cultivated on solid and in liquid media.

For Research Use Only. Not for use in clinical diagnostic procedures.

Please contact your local representative for availability in your country.

No. 1926931 © 02-2026 Bruker Daltonics

MALDI Biotyper® is a registered trademark of the Bruker group of companies.

## Online information

[bruker.com/microbiology](https://bruker.com/microbiology)



**Bruker Daltonics GmbH & Co. KG**

Bremen · Germany  
Phone +49 (0) 421-2205-0

[info.md@bruker.com](mailto:info.md@bruker.com)

**Bruker Scientific LLC**

Billerica, MA · USA  
Phone +1 (978) 663-3660

