



EDULAB FOR STUDENTS: MAGNETTECH ESR5000

TEMPO-tations in the Kitchen: Discovering Antioxidants in Olive Oil

EPR of Antioxidants in Olive Oil

Authors & Affiliation:

Reef Morse¹ and Kalina Rangelova²

¹Steppingstone MAgnetic Resonance Training Center, 650 Church Street, Plymouth, MI, USA

²Bruker BioSpin Corporation, 15 Fortune Drive, Billerica, MA, USA

Experiment Hashtag: #OliveOil, #Educate2Resonate

Keywords:

EPR, Olive Oil, Free Radicals, Antioxidants, Polyphenols, Food Analysis

Target Group:

Advanced Undergraduate or Graduate, General Chemistry, Analytical Chemistry, Food Chemistry, Food Safety and Control Laboratory, General Life Sciences

Objectives:

- Gain hands on experience with the Bruker Magnettech ESR5000 benchtop spectrometer while building upon the understanding of electron paramagnetic resonance (EPR) in a practical setting
- Apply EPR to a real-world sample
- Learn how to process and interpret EPR data and kinetics

Background of the Experiment:

The world we live in is a dangerous place for complex organisms. Radiation knocks the occasional molecule apart which produces reactive entities, called free radicals, that attach to other molecules in ways we might not want. Fortunately, over the last several billion years, our progenitors evolved mechanisms to shield themselves from these dangerous chemicals. This shield consists of a complex series of chemical reactions that metabolize these reactive molecules before they cause damage, repair damage if it happens, or outright capture and render harmless the reactive molecules directly. This last layer of protection uses molecules called antioxidants. Some organisms produce their own antioxidants while other get their antioxidants from what they eat.

It's useful to know if antioxidants are present in our food. We might prefer products that are high in these protective molecules if we have a choice. But food labels don't always give us that information. One antioxidant that is commonly listed is vitamin C, known to chemists and nutritionists as ascorbic acid. But much of the biochemistry that our bodies perform occurs in places where water, and hence vitamin C, is scarce. Our cells and their organelle are defined by cell or organelle membranes which are composed of lipids (a fancy term for fats). Membranes not only define our cellular systems but act as surfaces for a great deal of biochemical activity including the reactions that produce chemical energy by turning oxygen into water: a process that produces extremely reactive oxygen radicals (think of peroxide and superoxide). While we produce some antioxidants that protect our cells during these processes, we must get others from our food, and these antioxidants aren't part of the food labels.

The experiments described here that you will perform will give you a set of tools to measure antioxidant levels in food with high lipid content. We'll be studying olive oil but you can use this method on other foods as well.

We'll need some special equipment to do these experiments, but we will be able to get data we can't get any other way. The instrument we'll use, an EPR spectrometer, is purpose-built to measure molecules or atoms with unpaired electrons, as free radicals are. Because electrons are magnetic, they can be made to absorb energy in a magnetic field. How they return that energy to the sample can give us detailed information about the environment the electron is in. More importantly, for our purposes, because antioxidants attack free radicals, if we add free radicals to the sample and antioxidants are present, they will neutralize the free radical and make its signal go away.

Our experiment will therefore require a special kind of free radical; one that isn't particularly reactive but can be neutralized by antioxidants. All we need to do is keep track of the amount of this indicator molecule that is present in the sample that hasn't been neutralized. The indicator molecule we will use has the official chemical name of 2,2,6,6-tetramethylpiperidine-N-oxyl. We'll use its nickname TEMPO.




Figure 1 Chemical structure of 2,2,6,6-tetramethylpiperidine-N-oxyl (TEMPO)

Glossary

EPR: Electron paramagnetic resonance or electron spin resonance spectroscopy is a method for studying materials that have unpaired electrons. The basic concepts of EPR are analogous to those of nuclear magnetic resonance, but the spins excited are those of the electrons instead of the atomic nuclei.

FREE RADICALS: An atom, molecule, or ion that has at least one unpaired valence electron

ANTIOXIDANTS: Antioxidants are compounds that inhibit oxidation, a chemical reaction that can produce free radicals. Their scavenging activity is different. Some of the most potent antioxidants are vitamin E, vitamin C, phenol alcohols, etc.

If we make a very dilute solution of TEMPO in olive oil and put it in our EPR spectrometer, we'll see a signal like this:




Figure 2 EPR spectrum of TEMPO

The three lines come from the magnetic field of the electron interacting with the magnetic field of the nitrogen nucleus. The magnetic field applied to the sample increases from left to right. For reasons we won't go into here, the electron only absorbs energy at a specific magnetic field. This absorption happens at three different magnetic fields because the magnetic field of the nitrogen nucleus adds to, subtracts from, or has no effect on, the applied magnetic field from the instrument.

If we collect EPR spectra over time, we can watch the signal to determine how fast it disappears and from that intuit how fast the free TEMPO radical is being neutralized. The more antioxidant in our sample, the faster the signal will disappear.

In this exercise, we will analyze olive and soybean oil to determine their antioxidant properties. We will monitor TEMPO free radical signal being reduced by antioxidants in the two types of oil and calculate the rate of reduction.

Preparation:

To ensure efficient completion of the experiments, it is recommended to form groups with a maximum of three students. The estimated time for sample preparation is approximately 30 min. The EPR experiments are expected to take approximately 2.5 hours in total including two samples of oil — olive and soybean (the second oil is optional). After completing the experiments, an additional 1.5 - 2 hours will be needed to write a report. It is assumed that students have already covered introductory concepts of EPR and have a basic understanding of instrumental parameters.

For comprehensive information on EPR basics and optimizing instrumental parameters, students can consult the Magneotech ESR5000 educational kit, which is also provided with the benchtop EPR spectrometer.

To perform these experiments, an installed Magneotech ESR5000 spectrometer is required.

Experimental Setup:

Materials:

- 95 % ethanol
- TEMPO
- 1 mm x 100 mm glass capillaries
- 1.5 ml Eppendorf tubes
- 1 ml glass vial
- Pipettes to measure variable amounts of liquid from 2 μ l to 1 ml
- Balance accurate to at least 0.1 mg
- Capillary sealant
- Olive Oil (~ 5 ml)
- 2nd type of oil (optional)

Abbreviations

EPR: Electron Paramagnetic Resonance

References:

1. Ottaviani M. F. et al., *Electron paramagnetic resonance investigations of free radicals in extra virgin olive oils*, J. Agric. Food Chem. 49 (2001) 3691-3696
2. Lanza B. and Ninfali P., *Antioxidants in extra virgin olive oil and table olives: Connections between agriculture and processing for health choices*, Antioxidants 9 (2020) 41-58

Acknowledgments:

We wish to acknowledge the assistance of Emily, Nathan, Lucille, and Sophie Cheng of the SMART Center in testing the experimental details of the student portion of this laboratory.

